


- 16 Philip-Joet FF, Paganelli FF, Dutau HL, et al. Haemodynamic effects of bilevel nasal positive airway pressure ventilation in patients with heart failure. *Respiration* 1999;**66**:136–43.
- 17 Bradley T, Holloway R, McLaughlin P, et al. Cardiac output response to continuous positive airway pressure in congestive heart failure. *Am Rev Respir Dis* 1992;**145**:377–82.
- 18 Baratz DM, Westbrook PR, Shah PK, et al. Effect of nasal continuous positive airway pressure on cardiac output and oxygen delivery in patients with congestive heart failure. *Chest* 1992;**102**:1397–401.
- 19 Naughton MT, Rahman MA, Hara K, et al. Effect of continuous positive airway pressure on intrathoracic and left ventricular transmural pressures in patients with congestive heart failure. *Circulation* 1995;**91**:1725–31.
- 20 Pepin JL, Krieger J, Rodenstein D, et al. Effective compliance during the first 3 months of continuous positive airway pressure: a European prospective study of 121 patients. *Am J Respir Crit Care Med* 1999;**160**:1124–9.
- 21 Collard P, Pieters T, Aubert G, et al. Compliance with nasal CPAP in obstructive sleep apnea patients. *Sleep Med Rev* 1997;**1**:33–44.
- 22 Barbe F, Mayoralas LR, Duran J, et al. Treatment with continuous positive airway pressure is not effective in patients with sleep apnea but no daytime sleepiness: a randomized, controlled trial. *Ann Intern Med* 2001;**134**:1015–23.
- 23 Monasterio C, Vidal S, Duran J, et al. Effectiveness of continuous positive airway pressure in mild sleep apnea-hypopnea syndrome. *Am J Respir Crit Care Med* 2001;**164**:939–43.
- 24 Reeves-Hoche MK, Hudgel DW, Meck R, et al. Continuous versus bilevel positive airway pressure for obstructive sleep apnea. *Am J Respir Crit Care Med* 1995;**151**:443–9.
- 25 Hanly P, Zuberi-Khokhar N. Daytime sleepiness in patients with congestive heart failure and Cheyne-Stokes respiration. *Chest* 1995;**107**:952–8.
- 26 Kaneko Y, Floras JS, Usui K, et al. Cardiovascular effects of continuous positive airway pressure in patients with heart failure and obstructive sleep apnea. *N Engl J Med* 2003;**348**:1233–41.
- 27 Tkacova R, Liu PP, Naughton MT, et al. Effect of continuous positive airway pressure on mitral regurgitant fraction and atrial natriuretic peptide in patients with heart failure. *J Am Coll Cardiol* 1997;**30**:739–45.
- 28 Lenique F, Habis M, Lofaso F, et al. Ventilatory and haemodynamic effects of CPAP in left heart failure. *Am J Respir Crit Care Med* 1997;**155**:500–6.
- 29 Sanders M, Kern N. Obstructive sleep apnea treated by independently adjusted inspiratory and expiratory positive airway pressures via nasal mask: physiologic and clinical implications [see comments]. *Chest* 1990;**98**:317–24.
- 30 McAlister FA, Ezekowitz JA, Wiebe N, et al. Systematic review: cardiac resynchronization in patients with symptomatic heart failure. *Ann Intern Med* 2004;**141**:381–90.
- 31 Andreas S, Clemens C, Sandholzer H, et al. Improvement of exercise capacity with treatment of Cheyne-Stokes respiration in patients with congestive heart failure. *J Am Coll Cardiol* 1996;**27**:1486–90.
- 32 Granton J, Naughton M, Benard D, et al. CPAP improves inspiratory muscle strength in patients with heart failure and central sleep apnea. *Am J Respir Crit Care Med* 1996;**153**:277–82.
- 33 Ayas NT, Patel SR, Malhotra A, et al. Auto-titrating versus standard continuous positive airway pressure for the treatment of obstructive sleep apnea: results of a meta-analysis. *Sleep* 2004;**27**:249–53.
- 34 Kohnlein T, Welte T, Tan LB, et al. Assisted ventilation for heart failure patients with Cheyne-Stokes respiration. *Eur Respir J* 2002;**20**:934–41.

## IMAGES IN CARDIOLOGY .....

doi: 10.1136/hrt.2005.069872

### Angina from a right coronary artery to right atrial fistula

A 63 year old man with prior history of a motor vehicle accident and ensuing C5 quadriplegia presented with a diarrhoeal illness and dehydration. His other medical history was significant for paroxysmal atrial fibrillation, mild asthma, Addison's disease, and recurrent urinary tract infections. Prior echocardiography had demonstrated normal left ventricular function with a secundum atrial septal defect. During hospitalisation, he complained of frequent bouts of ischaemic chest pain. Electrocardiography revealed a right bundle branch block and ST segment depression in the inferior leads with pain. Despite anti-ischaemic treatment, his symptoms persisted and a coronary angiogram was performed. The left coronary system was angiographically normal. The dominant right coronary artery arose from the right coronary cusp and gave way to a highly tortuous and aneurysmal vessel that was well demonstrated during left ventriculography (panel A). A follow up transoesophageal echocardiogram confirmed the aneurysmal right coronary artery measuring 15 mm in diameter, with a fistulous communication into the right atrium (panels B–D). Left to right shunt was demonstrated from the right coronary artery into the right atrium but no atrial septal defect. He has thus far remained stable with only mild angina on effort and collaborative consideration is being given to how his anatomical anomaly would best be managed if symptoms were to intensify.



P Barlis  
P Calafiore  
D O'Donnell  
peter.barlis@gmail.com